




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61M 31/00	A1	(11) International Publication Number: WO 99/51296 (43) International Publication Date: 14 October 1999 (14.10.99)
(21) International Application Number: PCT/US99/06665 (22) International Filing Date: 26 March 1999 (26.03.99) (30) Priority Data: 60/080,566 3 April 1998 (03.04.98) US 09/087,146 29 May 1998 (29.05.98) US (71)(72) Applicant and Inventor: McDANIEL, David, H. [US/US]; Suite 113, 933 First Colonial Road, Virginia Beach, VA 23454 (US). (74) Agent: CHARTOVE, Alex; Akin, Gump, Strauss, Hauer & Feld, LLP, Suite 400, 1333 New Hampshire Avenue, N.W., Washington, DC 20036 (US).		(81) Designated States: AU, BR, CA, CN, IL, JP, KR, MX, NZ, SG, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>

(54) Title: ULTRASOUND ENHANCEMENT OF PERCUTANEOUS DRUG ABSORPTION



(57) Abstract

A system for enhancing and improving the transcutaneous or transdermal delivery of topical chemicals or drugs. A disposable container contains a substantially sterile unit dose of an active agent adapted for a single use in a medical treatment. The unit dose is formulated to enhance transport of the active agent through mammalian skin when the active agent is applied to the skin and the skin is exposed to light and/or ultrasound defined by at least one specific parameter.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LJ	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

ULTRASOUND ENHANCEMENT OF PERCUTANEOUS DRUG ABSORPTION

Related Cases

This application claims priority from co-pending provisional application serial no. 60/080,566, filed April 3, 1998.

Field of the Invention

The present invention generally relates to a system for enhancing and improving the transcutaneous or transdermal delivery of various topical chemicals or drugs (also referred to herein as "active agents").

Background of the Invention

Heretofore, ultrasound has been primarily used for diagnostic purposes with an outstanding safety record. It has also been used for dental care. Physical therapy uses ultrasound primarily to generate deep heat and also sometimes as an adjunct to wound healing. Some attempts have been made since the 1950's to use ultrasound to deliver hydrocortisone into joint spaces (originally for bursitis) or lidocaine (for pain relief) rather than injecting with needles. The current use of ultrasound to deliver drugs is primarily its use in physical therapy for non-invasive treatment of certain musculoskeletal disorders. Research in recent years has dramatically increased the understanding of ultrasound and its effects on skin and transport of topical agents. However, there is no consensus on how to optimally increase the flux or flow of topical agents across the skin using ultrasound.

Although ultrasound has been used to deliver drugs very deeply into joints, several problems exist with this technique. For example, if high frequency ultrasound is directed toward a bone for an extended period of time, then the energy can cause a burn. To some extent a focused beam tends to cause uneven concentration and uneven penetration, and may also cause injuries. This is particularly true with older ultrasound equipment, but is also true of many current ultrasound technology in clinical use today. An additional problem with

some earlier ultrasound equipment is that a two to four hour exposure period may be required. The newer ultrasounds use frequencies that provide results in a five to twenty minute time frame.

The use of ultrasound to deliver agents transcutaneously is generally termed "sonophoresis" but occasionally is termed "phonophoresis". Ultrasound generally comprises high-frequency sound waves that are above the human hearing range (usually greater than 20,000 Hertz (Hz) frequency units).

The sound waves may be generated by applying an alternating high frequency electrical current to a crystal such as a quartz, silicone dioxide, lithium sulfate of barium titanate. This current distorts the crystal, creating high frequency vibrations known as the piezoelectric effect. The sound waves produced have energy and may penetrate matter, depending on its acoustic density and composition. These sound waves may be delivered either in a focused manner and concentrated to a focal point (similar to the effects achieved with a magnifying lens and sunlight) or delivered in a non focused manner, termed "collimated," whereby the beam is uniform and parallel with no focal point (similar to falling rain).

The depth of penetration of ultrasound is inversely related to the frequency. Current diagnostic and therapeutic ultrasound typically ranges in frequency from 1-3 MHz to 4-10 MHz. Delivery may be pulsed in bursts or continuous beam modes, either stationary or continuously moving (usually at a rate of about one inch per second). The energies used generally range from a few milliwatts to a few watts.

The ultrasound energy is usually delivered through a transducer head. When used on skin, it is usually placed in direct contact with the skin using a coupling medium (which is often an aqueous gel), as shown, for example, in Figure 1.

It is also known that topical agents may be applied directly to the skin. Sometimes absorption of these agents may be enhanced by techniques such as

occlusion with synthetic materials (e.g. plastic), or transport may be enhanced by using low level electrical means such as galvanic current application or the technique of iontophoresis.

One problem with this known methodology is that of achieving more effective beneficial therapeutic clinical effects with topical agents by reaching higher concentrations in the dermis without damaging the dermis and/or the epidermis. Such damage may include undesirable and intolerable side effects such as flaking, peeling, persistent redness or burning.

The problem has not been adequately solved by simply using topical agents in various vehicles at higher strength. Using solutions at higher strength may tend to magnify the side effects. Using solutions at higher strength may also worsen internal systemic side effects. Simply using solutions at higher strength tends to work too broadly, and generally fails to selectively cause the desired effect.

Iontophoresis tends to work only if the molecules can be broken up into positive and negative ions and then driven in. A primary disadvantage of this technique is that many drugs cannot be broken up into positive and negative ions. Another disadvantage is that many drugs lose their effectiveness if broken up in this manner.

It has been proposed that drugs may be injected into the skin using a hypodermic needle. However, a primary disadvantage of this technique is that it generally fails to deliver the drug uniformly.

It has been proposed that drugs may be taken orally. However, in many cases this technique is not practical. For example, oral ingestion of vitamin C generally fails to provide a sufficiently high concentration of the drug in the skin. The technique may also involve certain risks. For example, oral ingestion of vitamin A derivatives may cause liver damage or birth defects.

It can be seen, therefore, that while the drugs themselves may be quite effective if used selectively, each of the traditional routes has one or more

inherent limiting factors which prevent or inhibit the drugs from achieving their maximum effectiveness.

Summary of the Invention

In accordance with the present invention, these and other objectives may be achieved by providing an optimal selection of ultrasound parameters such as frequency, intensity, pulse length, beam characteristics and application time on the skin (including both human and animal) to enhance the transport of topical agents into epidermal, dermal and subcutaneous tissues. The present invention may be used to produce higher concentrations of such an agent than may be accomplished by current topical application or delivery methods. Such increased concentrations may have beneficial effects depending on the characteristics of the topical agent delivered. Patients who cannot tolerate current topical application methods may achieve beneficial effects by delivering similar concentrations of the agent with little or no side effects. Topical agents which cannot normally penetrate the skin with current methods may be transported into the dermis

The effects of ultrasound on skin result from the release of energy. These include the non-thermal effects of (1) cavitation, (2) mechanical stress as well as (3) thermal effects.

It appears that ultrasound exposure in the therapeutic range causes cavitation in the keratinocytes of the stratum corneum as the primary effect in increasing skin permeability for transcutaneous transport of topical agents (cavitation is a process where bubbles are formed which oscillate causing structural disorder of the intercellular lipid bilayers of the keratinocytes).

In effect, this process is somewhat analogous to loosening the "mortar between the bricks" and expanding the "spaces between these bricks" so that the topical agent has a transport pathway to reach the epidermis and the deeper dermis of the skin. Drug molecules that are too large to penetrate the skin at all

when applied topically may achieve significant penetration when used in conjunction with sonophoresis with proper parameters. See Figure 3 and 4.

Brief Description of the Drawings

A detailed description of a preferred embodiment of the present invention will be made with reference to the accompanying drawings.

Figure 1 illustrates an example of the delivery of ultrasound energy through a coupling medium and a transducer head.

Figure 2 illustrates an example of the percutaneous delivery of topical agents being principally limited by the barrier function of the stratum corneum (the outermost 15-25 microns of the skin).

Figures 3 and 4 illustrate an example of cavitation from ultrasound "expanding" the spaces between keratinocytes, thus creating a channel for drugs including drug molecules that are too large to penetrate the skin when applied topically achieving significant penetration when used in conjunction with sonophoresis with proper parameters.

Figure 1.1 illustrates an example of a pretreatment of the stratum corneum.

Figure 1.2 illustrates an example of a coupling agent being applied and an ultrasound treatment being performed.

Figure 1.3 illustrates an example of a coupling agent being removed and an active topical agent being applied.

Figure 1.4 illustrates an example of an active topical agent being removed after an appropriate time.

Figure 1.5 illustrates an example of a protective topical agent being applied.

Figure II.2 illustrates an example of an active topical agent which also serves as a coupling agent being applied, and an ultrasound treatment being performed.

Figure III.1 illustrates an example of an injection of a drug or agent or autologous or donor fibroblast rich preparation.

Figure IV illustrates an example of an embodiment using low energy laser therapy to stimulate (directly or indirectly) the production, proliferation, activation, or inhibition of the activity, structure or function of various biochemical or photochemical or biological processes (this example features stimulation of a human fibroblast).

Detailed Description of the Preferred Embodiments

The following detailed description is of the best presently contemplated mode of carrying out the invention. This description is not to be taken in a limiting sense, but is made merely for the purpose of illustrating the general principles of the invention. The scope of the invention is best defined by the appended claims.

In a preferred embodiment, each active agent may be specially formulated to achieve a desired result using ultrasound. At present, many drugs are formulated to achieve a desired result using topical penetration, but the inventor is unaware of any drug that is specially formulated to achieve a desired or optimal result using ultrasound. At present, a coupling agent must typically be used to enable effective transmission using ultrasound.

In a preferred embodiment each active agent may be provided in a single dose container or packet. The packet may preferably contain only a pure active agent, at an optimal strength and/or at an optimal concentration for a specific use. The active agent may be dissolved in an appropriate vehicle medium, such as a gel or a liquid. All other contaminants (including, for example, preservatives, fragrances, etc.) are preferably absent or removed from the packet. The need to include, with the active agent, additives for prolonging shelf life, improving stability, preventing spoilage, improving texture, etc., is thereby eliminated. In a preferred embodiment of the invention everything is removed

from the active agent formulation which does not have a desired effect. Each packet is preferably packaged in a sterile manner, by radiation sterilization or other means. Additionally, the parameters of the collimated ultrasound are preferably selected for maximum safety. Preferably, exposure times are confined to intervals of five, ten, fifteen or twenty minutes.

In this preferred embodiment a layer of the active agent may first be placed on the skin. The ultrasound equipment may be adjusted to optimize the ultrasound parameters, including the exposure time, and then turned on. The ultrasound wand may be moved gently over the skin for the duration of the selected exposure time. The treatment is essentially complete at the end of the selected exposure time. Any active agent residue left on the skin may be cleansed with a gentle cleansing agent, so that no active agent is left to irritate the skin. Treatment may be repeated at periodic intervals.

In another embodiment of the invention an ultrasound gel or similar substance may be put on the skin first. The skin may then be exposed to ultrasound. The gel may be removed and the active agent may be placed on the skin. The active agent may be allowed to sit on the skin for a period of time, preferably between about five and twenty minutes. This tends to provide a short "burst" of the active agent at a higher penetration. The residue may then be wiped off, so that no active agent is left to irritate the skin.

Another option would be to apply the active agent in a 'delivery device' such as a biomembrane, dressing or band-aid, liposome-polymer complex or aerosol spray dressing, which functions as a delivery device, but also provides a barrier function. In this option, the active agent would be left on the skin for a period of hours or perhaps overnight.

It appears that, at least in this embodiment, the increased permeability or barrier spreading on the cells may last for up to one or two days. A topical agent, aerosol agent, dressing, or delivery device may be applied to help protect during this time or to help restore the barrier function. A preferred formulation

could be a lipid mixture such as cholesterol, ceramides, free fatty acids, linoleic acid. Another option would be a lipid or liposome - polymer mixture or aerosol polymer coating. Consequently, the skin may be relatively absorbent and it may be preferable to provide some associated aftercare, such as avoiding sunburn, Retin-a or aftershave. Treatment with a topical agent specifically formulated to "restore" or improve barrier function may be used. Such a topical agent may also be packaged as described above.

In another embodiment of the invention an outer layer of skin may be removed before the skin is exposed to an active agent or ultrasound. This tends to enable the active agent to penetrate better and/or deeper. This may be achieved, for example, by first wiping the skin with acetone to strip the oils out. An enzyme may then be used to selectively kill only dead skin cells. Pretreatment with heat, skin hydrating preparations and preparations to alter (and optimize for treatment) the skin pH may also be utilized. Chemicals, abrasers and lasers may also be used for this purpose, although they tend to be less discriminate. An active agent may then be placed on the skin and exposed to ultrasound, or the skin may be exposed to ultrasound prior to placing an active agent on the skin. The active agent may then be wiped off. The barrier function of the outer skin layer tends to return within a few hours or days.

It has been found that removing impediments to the introduction of an active agent to the skin tends to have at least one of the following three effects: (1) increasing the amount of active agent which enters the skin, (2) shortening the time during which the skin must be exposed, and/or (3) increasing the penetration depth of the active agent. Precision, uniformity and safety may be improved by reducing variations in the material actually placed on the skin and having the ultrasound beam as uniformly distributed as possible.

Treatment in accordance with the present invention tends to be relatively safe, relatively inexpensive, does not necessarily need to be performed by a physician, has a recovery time which is relatively short (and in many cases non-

existent), and can be done repeatedly with virtually no cumulative adverse effects but with cumulative beneficial effects.

The present invention may provide a safe, effective delivery mechanism for a wide variety of active agents, including bleaching cremes, vitamin C, the vitamin A family, and topical anesthetics. In some cases the present invention may allow topical anesthetics to be administered in sufficient strength that scalpel work may be performed without needles. The present invention may be used for the treatment of leg ulcers, where the known effects of ultrasound alone may be used to stimulate wound healing and treatment in accordance with the present invention may be used to drive one or more new agents which enhance, speed or stimulate wound healing in to greater depths, producing results that are better and/or faster.

The packets may be provided in different sizes. For example, the size of one packet may be configured so that there is sufficient active agent in the packet to treat the average face. As another example, packet size may be configured so that there is sufficient active agent in the packet to treat the arms and/or legs. The packet may be configured for unit dose delivery, and the active agent may be optimized both for its vehicle and its percentage.

If two or three active agents are used in combination, or if some of the active agents are dissolved in micro spheres or synthetic devices, then the penetration ratio may be affected by the coefficients of diffusion and characteristics that are unique for each active agent, including molecular size and shape. If two or more active agents are used in combination, then each active agent may not necessarily be driven into the skin at the same ratio because each molecule may be driven in at a different rate. When several active agents are used in combination, some active agents may be driven in to higher strengths than others based on factors including their weight, their volume, their size, their shape and other properties. These factors may be optimized to achieve the desired treatment effect.

The active agent may be used to stimulate biological systems, and from a disease therapeutic standpoint the active agent may also be used to inhibit or turn off certain biological functions. In other words, the present invention may be used to provide an active agent concentration of sufficient magnitude that it may cause a feedback inhibition. For example, if a concentration of 2% in the skin provides stimulation and a concentration of 4% is considered optimal, then it is possible that a concentration of 10% may provide less effect than 4% and at 15% the reverse effect may occur. It is therefore possible to have an active agent formulation (and/or time period and/or set of parameters) optimized so that the treatment does not merely restore biological function but actually treats disease.

The embodiment illustrated in Figure IV is primarily as a 'biostimulating' process and not a destructive or ablative process, such as commonly used in today's art. Using laser therapy or ultrasound (or both) to produce heat at a subnecrotic damage or injury threshold is another possible embodiment. (An example might be trying to achieve a dermal temperature that affects collagen - perhaps beginning around 63-65°C).

Figure IV illustrates an example of an embodiment using low energy laser therapy to stimulate (directly or indirectly) the production, proliferation, activation, or inhibition of the activity, structure or function of various biochemical or photochemical or biological processes so as to result (directly or indirectly) in effects that are beneficial to the structure, function or appearance of the skin and/or subcutaneous tissues or which results in the "rejuvenation" of photoaged, environmentally damaged or disease or drug/therapy altered skin or subcutaneous tissue.

The present invention, in a preferred embodiment, may be used either as a pretreatment followed by topical application of a desired agent or used in a coupling media vehicle to directly deliver the desired agent. At present, topical

pharmaceutical products are not generally formulated in such a way as to optimize their efficiency as ultrasound couplants.

A preferred formulation for an "active" coupling agent is one which would best meet the criteria described for the optimal active agent alone, and yet would still provide an effective coupling function for an ultrasound transducer head.

The present invention, in a preferred embodiment, formulates these topical agents in concentrations and vehicles so as to be optimally effective when used in conjunction with sonophoresis. These specifications may vary with molecular size as well as diffusion and partition coefficients and other factors and may be uniquely formulated for each agent. The preferred embodiment for many may be a gel. Also preferred is individual single use sterile dose packets, thereby avoiding the need for preservatives and other additives that may cause undesirable effects or side effects.

The goal in many cases may be to maximize percutaneous transmission of the active agent. The following principles may be useful in selecting these formulations, except in some situations where a milder or submaximal effect or transmission may be desired for a given agent or situation. In such case, the formulation may be shifted or modified to have reduced effect. Also, certain complex or synthetic delivery systems may follow different principles, such as lipid bilayers (liposomes) or synthetic polymers (microsponges).

However, for many situations the preferred formulation may be based upon the desire to approach, but not quite reach, a saturated solution of the active agent(s). Thus, the rate of release of active agent or drug may be greater in vehicles in which it exhibits poorest solubility. This may be more important than the total concentration, so as to maximize bioavailability and release.

In a preferred formulation, it may be desirable to select a vehicle which poorly solubilizes the active agent and which preferably may evaporate at such a rate that the concentration of the active agent increases to offset the release into and through the stratum corneum. However, it may also be important that it not

evaporate so quickly as to precipitate out on the skin, since this may decrease or even stop transport (situations where stable, supersaturated solutions form may be acceptable). Thus, it may be desirable to have sufficient non-volatile vehicle to avoid precipitation from occurring during the time the active agent may be applied to the skin. Since the transport rate may be affected by various factors, such as those described herein, the optimal vehicle may vary not only with each specific active agent, but also with the specific technique parameters (or disease state) since the barrier function (and thus transport depletion of concentration of active agent in vehicle) may vary.

Preferred ultrasound parameters may be determined in accordance with both efficacy and safety requirements. For example, one preferred range for lower frequency ultrasound may be between about 25 kHz and about 3 MHz at about 0.5 - 2.0 W/cm² (either continuous or pulsed, using about a 20 - 25% duty cycle if pulsed). A preferred setting within this range may be at about 1.0 MHz at about 2.0 W/cm², with a continuous wave beam and a treatment time of between about five and ten minutes. A preferred setting within this range for non-continuous beam (i.e. pulsed delivery) may be at about 1.0 MHz at about 0.2-0.5 W/cm² with a 20-25% duty cycle with around a 2.0-20.0 msec "on" cycle and a treatment time of about five and ten minutes. One preferred range for higher frequency ultrasound may be between about 3 MHz and about 16 MHz at about 0.2-1.0 W/cm² (either continuous or pulsed, using about a 20% duty cycle if pulses), with a treatment time of between about one and twenty minutes. A preferred setting within this range may be at about 10 MHz at about 0.2 W/cm², with a continuous treatment time of between about five and twenty minutes.

In a preferred embodiment the beam profile may be a collimated beam with precise control of output. In an alternative embodiment a focused beam may also be suitable. A preferred embodiment may provide a feedback warning if the operator loses contact with the skin, thus ensuring optimal treatment technique.

Further adjuncts to the process which increase permeability of skin or decrease skin barrier function may be helpful with optimizing the present invention. Options for this include, but are not limited to, stripping, removing, thinning or diminishing the structure, function, thickness or permeability of the stratum corneum by various mechanical, abrasive, photo acoustical, ablative, thermal, chemical, abrasive or enzymatic methods. Examples of these could include solvent or tape stripping, scrubbing, laser ablation or vaporization, chemical peeling, micro dermabrasion, enzyme peeling, or laser treatment using high peak power, short pulse duration lasers. A preferred embodiment may be enzyme peel, which is formulated to specifically remove only the dead stratum corneum cells.

In another embodiment of the invention sonophoresis may be used alone, without a topical agent, to produce a thermal effect (rather than to drive drugs in) to stimulate the skin (e.g. make fibroblasts, produce new collagen, elastin, etc.).

Examples of the active agent might include any of the following, either alone or in combination: Vitamin C; Vitamin E; Vitamin A; Vitamin K; Vitamin F; any of the various chemical forms and analogs of these vitamins; Retin A (Tretinoin); Adapalene; Retinol; Hydroquinone; Kojic acid; various growth factors; echinacea; antibiotics; antifungals; antivirals; bleaching agents; alpha hydroxy acids; beta hydroxy acids; salicylic acid; antioxidant triad compound (with or without Tretinoin or Vitamin A derivatives); seaweed and salt water derived products antioxidants, phytoanthocyanins, phytonutrients, botanical and herbaceous products, hormones (including insulin or estrogens), enzymes, minerals, growth factors, genetically engineered substances, cofactors or catalysts for various biological pathways and other antiaging substances.

The active agents used in accordance with the present invention may be characterized by one or more of a variety of properties. Optimization of the active agents for use in accordance with the present invention may be achieved by the modifying one or more of these properties. The following are examples of

some of the properties which may be modified, alone or in combination, to optimize active agents for use in accordance with the present invention:

Solubility - A sufficient concentration of active agent should preferably be dissolved in the selected delivery system vehicle or coupling agent for the desired treatment.

Stability - Some active agents may be unstable and may rapidly degrade after being dissolved in a vehicle or coupling agent. Therefore, in one preferred embodiment the active agent may be in the form of a powder (such as a freeze dried powder or lyophilized powder, for example) that is not mixed until treatment is imminent. Other instabilities may be related to oxidation from atmospheric oxygen or exposure to ultraviolet light or sunlight. Thus, in one preferred embodiment the active agent may be packaged under vacuum or nitrogen or another inert gas and may be packaged in a manner that protects the active agent from light.

Molecular Size - The active agent should preferably have a molecular size which enables it to penetrate the skin at the time of maximum permeability, and then should preferably be in a form which is either "active" or may become "activated" in the skin. For example, in the case of Vitamin C treatments, L-ascorbic acid has stability problems but is the active form in the skin and is a small molecule which enables penetration. In contrast, magnesium ascorbyl phosphate is very stable but is a much larger molecule and does not penetrate easily with current delivery systems. Guy & Potts have shown that the permeability of human (and mouse) stratum corneum may be determined by the molecular volume (weight) and the partition coefficient log poor.

pH - The activity of an active agent may frequently depend on an appropriate pH level. It is also important for skin to be at an optimal pH level. There is good evidence that the stratum corneum is much more permeable to neutral molecules than the salts of the weak acids or bases. Furthermore, skin enzymes and other processes may operate optimally at certain pH levels and

poorly at others. For example, an enzyme preparation used to remove stratum corneum may operate effectively at a pH level of 8.5. The skin may preferably be "pretreated" with a topical agent that increases the pH of the skin to this level. At the end of the treatment it may be advantageous to add another topical buffering solution or agent to "readjust" the skin back down to a normal skin pH level of 5.5.

pKa - The pKa tends to affect the amount of free acid and base in equilibrium, and thus has an impact on both the beneficial and adverse clinical effects of the active agent. A recent study by the FDA has shown that there may be considerable variation in the beneficial clinical effects (as well as the adverse and undesirable clinical side effects) of glycolic acid, even when the glycolic acid concentrations are the identical percentage. For example, if the pH of the glycolic acid preparation is adjusted to around 4.4 (approximately equal to the pKa of glycolic acid), then the ratio of beneficial to adverse side effects is much better than when the pH is lower than the pKa (and more free acid is available in the equilibrium). The pKa may vary for different compounds and thus may effect the choice of formulation.

Purity, Sterility, Absence of Nonactive Ingredients - The presence of an impurity (including chemical contaminants, preservatives, additives, fragrances, or micro-organisms, for example) may also potentially have enhanced penetration and thus produce undesired adverse or toxic effects.

Lipid Bi-Layer and Ionization - These may determine the "distribution" of the active agent, either in the lipid phase or in the ionized chemical form. These may affect diffusion and delivery of the active agent.

Lipophilic and Lipophobic - The choice of the delivery vehicle or coupling agent for the active drug may be effected by these properties of the active agent.

Diffusion Coefficient - One of the dominant factors in determining the transfer of the active agent into the skin, the Diffusion Coefficient measures the use of movement across the stratum corneum and the difference in

concentration or percentage of active agent and also the thickness of the stratum corneum.

Partition Coefficient - One of the dominant factors in determining the percentage transfer of the active agent into the skin, the Partition Coefficient refers to the tendency for the active agent to leave its vehicle and enter the stratum corneum. The amount of material that moves across the units representative of stratum corneum and a given unit of time tends to be directly proportional to the Partition Coefficient as well as the Diffusion Coefficient.

In a preferred embodiment the invention may be provided in the form of a "treatment kit" prepackaged to contain all the appropriate topical agents, drugs, supplies, etc. needed for a specific treatment. The kit may be divided into compartments or zones. Each compartment may be coded by color, number or letter (for example), so that the compartments follow a "Step System" that guides the treatment provider (i.e., the nurse, esthetician, physician, etc.) through the treatment process. Within each color coded zone, individual products that are color coded to match that step color would be provided so that they would not be easily confused.

Each kit may be produced specifically for a certain active agent. Each kit may be produced specifically for a certain usage, based on disorder being treated, anatomic area, etc.

Treatment Regimen Cards may be included in the kit. The Treatment Cards may become a part of the patient record or chart. The Treatment Cards may enable the treatment provider to chart multiple, serial treatments. Such Treatment Cards may be used to record basic information such as date of treatment, anatomic location, disorder, patient name, comments, etc. The Treatment Cards may also provide information regarding suggested treatment intervals, numbers of treatments, incremental increases in treatment parameters, incremental increases in active agent application time or strength.

Several different Treatment Cards may be included in a kit for a treatment provider to select. For example, a single kit may include different Treatment Cards for sensitive skin, regular skin, and post surgical skin treatment. These different Treatment Cards may have, as their distinguishing characteristics, differences in one or more of at least the following: percentage dilution of active agents, amount of time various agents or treatment are applied, or ultrasound or light source parameters, etc. Different Treatment Cards may be provided for different anatomic areas, such as the face, the arms or the legs, where these parameters may also be varied. Different Treatment Cards may be provided for different treatment options, for example gentle treatment for mild sun damage vs. aggressive treatment for severe sun damage.

The Treatment Cards may be color coded to indicate the time period for equipment parameters directly on the card. The Treatment Cards may also cross reference for percent concentration to the kit itself (in essence within each step of the kit) which may be color coded (for example pale blue background). For example, there may be three levels of dilution or strength (mild, intermediate, maximal) which may be identified by a color code on the Treatment Card.

For example, at one step of the treatment the active agent may be mixed from a dry powder. The mixing instructions may be imprinted on the box for the dilution, and the Treatment Card may also indicate the recommended dilution. For example, a patient with sensitive skin receiving a series of six treatments may start at the most mild of the three dilution strengths for the first three treatments and then go to the intermediate for the last three. In contrast, a person with average skin might have the first two treatments performed at the mild strength, the third and fourth treatments at the intermediate strength, and the fifth and sixth treatments at the maximum strength. In that case weeks one and two of the Treatment Card may be one color to match the mild bar and the one background color, weeks three and four may be a different color to match

the intermediate bar, and weeks five and six may be a third background color to match the maximum bar.

Another option may be to have colors, letters or numbers (for example 1.0, 2.0, 3.0, etc.) for the strengths of the kits where a dilution was not involved (for example cream or lotion that does not require mixing from a powder). In such a case, the patient Treatment Card background colors may correspond to the strength or color of the kit, rather than an individual dilution or compound of the kit. In such a case the cream may be in separate packages outside the kit, but still color coded. Likewise, different color bars may be used to indicate the duration of the treatment in minutes, or the treatment parameters to be used on the ultrasound or laser. Colors may be also used in color code to link or "intertwine" ultrasound/topical treatments and light or laser light treatments into a predetermined sequence and at the appropriate time interval.

In a preferred embodiment of the invention, for example, Step One of the kit may include the following items:

- A tear and wipe prepackaged skin pad impregnated with a skin cleanser.
- A sterile unpreserved unit dose package, vial, twist off plastic, tear packet, etc. type of container which contains a solution that adjusts the pH of the skin to pH 8.5
- A vial of dry enzyme powder (which requires pH 8.5 to properly function).
- A small vial of an appropriate amount of sterile distilled unpreserved H₂O to add to the dry powder.
- A small plastic cup and plastic spatula to mix these in.

All of the items in Step One of the kit may be coded with the same identifying number, letter and/or color.

Step Two of the kit may include a set of instructions to use the coupling gel and ultrasound as per the patient Treatment Card parameters.

Step Three may include the active agent itself, in a sterile single use package, with any additional ancillary items needed for that particular agent.

Step 4 may include the post treatment skin protective regimen which may include the following:

- A cleansing agent to remove the specific active agent prepackaged.
- A buffering solution to restore the skin to pH 5.5 (if necessary for a given process or active agent).
- An environmental protection lotion which may be specially formulated to help protect the skin until it has regained its own barrier function.
- A possible nonirritating sun block to use on top of the environmental protective cream.

The present invention has numerous potential beneficial uses. For example, the present invention, in a preferred embodiment, may be used to directly or indirectly produce beneficial effects by biochemical stimulation of tissue or cells (or thermal events without the use of topical agents) including such activities as stimulating fibroblast production of new collagen or elastic fibers. The present invention, in a preferred embodiment, may be used to lighten uneven or extra pigment (including melanin) in the epidermis or dermis. The present invention may be used to add, augment or supplement antioxidants, vitamins, phytonutrients, trace elements, minerals, or naturally occurring, synthetic or generically engineered substances which may alter or improve the structure, health or function of the skin or subcutaneous tissue.

The present invention, in a preferred embodiment, may be used to: deliver agents which enhance, speed or promote wound healing; deliver anesthetic agents to the skin and subcutaneous tissues; improve skin tone and "tighten" loose skin; to reduce the appearance of cellulite; reduce wrinkles or scars; deliver drugs for the purpose of producing a non-local, systemic effect (such as insulin); deliver agents in their pure or neat form or in vehicles such as gels, creams, ointments, emulsions, micropolymer beads (or sponges, etc.), liposomes or other natural or synthetic "transport device" used with or without a coupling media.

The present invention provides the possibility for the non-surgical rejuvenation of skin, improved wound healing, better anesthetic for cutaneous surgery without needles, delivery of drugs that currently are ineffective or poorly tolerated by topical delivery.

The presently disclosed embodiments are to be considered in all respects as illustrative and not restrictive, the scope of the invention being indicated by the appended claims, rather than the foregoing description, and all changes which come within the meaning and range of equivalency of the claims are therefore intended to be embraced therein.

What is claimed is:

1. A method for enhancing the transport of an active agent through mammalian skin, comprising:
 exposing the skin to ultrasound, and
 applying an active agent to the skin.
2. The method of claim 1, wherein the step of exposing the skin to ultrasound and the step of applying an active agent to the skin occur substantially simultaneously.
3. The method of claim 1, wherein the step of applying an active agent to the skin occurs substantially after the step of exposing the skin to ultrasound.
4. The method of claim 1, wherein the step of applying an active agent to the skin comprises applying the active agent to the skin topically.
5. The method of claim 1, wherein the step of applying an active agent to the skin comprises injecting the active agent into the skin.
6. The method of claim 1, wherein the skin comprises a stratum corneum and wherein the method comprises the step of modifying at least a portion of the stratum corneum substantially prior to the step of exposing the skin to ultrasound.

7. The method of claim 6, wherein the step of modifying at least a portion of the stratum corneum comprises at least one of the steps of:

removing at least a follicular plug from the stratum corneum,
removing at least skin debris from the stratum corneum, and
removing at least a portion of the stratum corneum.

8. The method of claim 1, comprising the step of providing a coupling agent for enabling transmission of ultrasound through the skin.

9. The method of claim 8, wherein the step of applying an active agent to the skin comprises applying the active agent to the skin topically, and wherein the step of providing the coupling agent and the step of applying the active agent to the skin topically occur substantially simultaneously.

10. The method of claim 9, wherein the active agent and the coupling agent are substantially identical.

11. The method of claim 1, comprising the step of applying a protective agent to the skin.

12. The method of claim 1, comprising the step of exposing the skin to low energy light.

13. The method of claim 12, wherein the step of exposing the skin to low energy light occurs substantially prior to the step of exposing the skin to ultrasound.

14. The method of claim 12, wherein the step of exposing the skin to low energy light occurs substantially after the step of exposing the skin to ultrasound.

15. The method of claim 12, wherein the step of exposing the skin to low energy light occurs substantially prior to the step of applying an active agent to the skin.

16. The method of claim 12, wherein the step of exposing the skin to low energy light occurs substantially after the step of applying an active agent to the skin.

17. The method of claim 5, wherein the active agent comprises human fibroblasts.

18. The method of claim 12, wherein the step of applying an active agent to the skin comprises injecting the active agent into the skin.

19. The method of claim 18, wherein the active agent comprises human fibroblasts.

20. The method of claim 1, wherein the active agent comprises at least one of: Vitamin C; Vitamin E; Vitamin A; Vitamin K; Vitamin F; Retin A (Tretinoin); Adapalene; Retinol; Hydroquinone; Kojic acid; a growth factor; echinacea; an antibiotic; an antifungal; an antiviral; a bleaching agent; alpha hydroxy acid; beta hydroxy acid; salicylic acid; antioxidant triad compound; seaweed; a salt water derived product; an antioxidant, a phytoanthocyanin, a phytonutrient, a botanical product, a herbaceous product, a hormone, an enzyme, a mineral, a growth factor, a genetically engineered substance, a cofactor for a biological pathway, a catalyst for a biological pathway, and an antiaging substance.

21. A disposable container adapted for a single use and containing a substantially sterile, unit dose of an active agent for use in a single medical treatment, the unit dose being specifically formulated to optimize a specific medical effect when used in combination with at least one specific ultrasound parameter.

22. A disposable container containing a substantially sterile unit dose of an active agent adapted for a single use in a medical treatment, the unit dose being formulated to enhance or maximize transport of the active agent through mammalian skin when the active agent is applied to the skin and the skin is exposed to ultrasound defined by at least one specific ultrasound parameter.

23. A disposable container containing a substantially sterile unit dose of an active agent adapted for a single use in a medical treatment, the unit dose being formulated to enhance or maximize transport of the active agent through mammalian skin when the active agent is applied to the skin and the skin is exposed to light defined by at least one specific light parameter.

24. The method of claim 1, wherein the step of applying an active agent to the skin comprises applying at least one of a dressing, a patch, and a biomembrane to the skin.

25. The method of claim 5, wherein the active agent comprises a preparation containing human fibroblasts.

26. A kit comprising:

at least one disposable container containing a substantially sterile unit dose of an active agent adapted for a single use in a medical treatment, the unit dose being formulated to enhance or maximize transport of the active agent through mammalian skin when the active agent is applied to the skin and the skin is exposed to at least one of ultrasound defined by at least one specific ultrasound parameter and light defined by at least one specific light parameter,

at least one substance adapted to be applied to the skin before the active agent is applied to the skin, and

at least one substance adapted to be applied to the skin after the active agent is applied to the skin.

1 / 7

Figure 1





Figure 2



2 / 7

Figure 3

"Normal"





Figure 4

"With Cavitation Bubbles"



3 / 7




Figure 5




Figure 5a




Figure 5b




Figure 5c




Figure 5d






Figure 5e

4 / 7






Figure 8d




Figure 8c




Figure 8b




Figure 8a

6 / 7




Figure 9

7 / 7

Pigment Lightening Treatment - Facial Skin - Sensitive				
Initial TX Step 1	Week 2	Week 4	Week 6	Week 8
Enzyme 5min	Enzyme 10min	Enzyme 10min	Enzyme 10min	Enzyme 10min
Step 2				
Ultrasound 0.35W 5 min	0.35W 7 min	0.35W 10 min	0.50W 10 min	0.50W 10 min
Step 3 Skin Lightening Agent 1.0/5min				
Step 4	1.0/10 min	2.0/10 min	2.0/10 min	2.0/10 min

Figure 10

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61M 31/00

US CL : 604/501

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 604/20-22, 500-502, 206, 511, 890.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS

Search Terms: (ultrasound (p) skin (p) drug)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5,636,632 A (BOMMANNAN et al.) 10 June 1997, Abstract.	1-26



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents.	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance, the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance, the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

09 MAY 1999

Date of mailing of the international search report

04 JUN 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

DEBORAH BLYVEIS

Telephone No. (703) 308-2110